

UCSF Health Hyde Hospital Antibigram 2025

| Percent Susceptible 2025 Isolates  | | Number of Isolates | Aminoglycoside | β-Lactam | | Beta lactam inhibitor combo | | Cephalosporin | | | Carbapenem | | Folate pathway | Fluoroquinolone | | Glycopeptide | Lincosamide | Tetracycline | Oxazolidinone |
|--|-------------------------------------|--------------------|------------------------------|------------|---------------------|-----------------------------|-------------------------|---------------|------|------|------------|-----------|-------------------------------|-----------------|--------------|--------------|-------------|--------------|---------------|
| | | | Gentamicin | Ampicillin | Oxacillin/Nafcillin | Ampicillin/sulbactam | Piperacillin-tazobactam | 1st | 3rd | 4th | Ertapenem | Meropenem | Trimethoprim/sulfamethoxazole | Ciprofloxacin | Levofloxacin | Vancomycin | Clindamycin | Tetracycline | Linezolid |
| | | | | | | | | | | | | | | | | | | | |
| Gram Negative | <i>Acinetobacter baumannii</i> | 9 | 100% | - | - | 100% | - | - | 78% | 100% | - | 100% | 89% | 100% | 100% | - | - | - | - |
| | <i>Citrobacter sp</i> | 7 | 100% | - | - | - | 83% | - | 85% | 100% | 100% | 100% | 100% | 100% | 100% | - | - | - | - |
| | <i>Enterobacter cloacae</i> | 28 | 100% | - | - | - | 75% | - | 71% | 86% | 82% | 96% | 89% | 96% | 96% | - | - | - | - |
| | <i>Escherichia coli (non-esbl)</i> | 177 | 91% | 54% | - | 62% | 94% | 79% | 100% | 100% | 100% | 100% | 71% | 72% | 79% | - | - | - | - |
| | <i>E. coli-ESBL (20% of total)</i> | 43 | 79% | - | - | - | - | - | - | - | 100% | 100% | 30% | 7% | 26% | - | - | - | - |
| | <i>Pseudomonas aeruginosa</i> | 69 | - | - | - | - | 80% | - | - | 81% | - | 87% | - | 74% | 69% | - | - | - | - |
| | <i>Klebsiella pneumoniae</i> | 55 | 96% | - | - | 76% | 93% | 89% | 95% | 95% | 96% | 100% | 89% | 89% | 91% | - | - | - | - |
| | <i>Proteus mirabilis</i> | 43 | 84% | 60% | - | 72% | 97% | 16% | 91% | 91% | 100% | - | 74% | 79% | 86% | - | - | - | - |
| | <i>Stenotrophomonas maltophilia</i> | 4 | - | - | - | - | - | - | - | - | - | - | 100% | - | 100% | - | - | - | - |
| Gram Positive | <i>Enterococcus faecium</i> | 9 | - | 11% | - | - | - | - | - | - | - | - | - | - | - | 56% | - | 22% | 100% |
| | <i>Enterococcus faecalis</i> | 20 | - | 100% | - | - | - | - | - | - | - | - | - | - | - | 55% | - | 15% | 100% |
| | <i>MR-S. aureus (52% of total)</i> | 119 | - | - | 0% | - | - | - | - | - | - | - | 90% | - | - | 100% | 74% | 54% | 100% |
| | <i>MSSA Staphylococcus aureus</i> | 109 | - | - | 100% | - | - | 100% | - | - | - | - | 88% | - | - | 100% | 85% | 94% | 100% |
| | <i>Coagulase Neg Staph</i> | 24 | - | - | 71% | 71% | - | 71% | - | - | - | - | - | - | 88% | 100% | 79% | 75% | 100% |
| | <i>Strep pneumoniae</i> | 13 | Penicillin-Susceptible: 100% | | | | | | | | | | | | | | | | |