

ADULT ANTIMICROBIAL SURGICAL PROPHYLAXIS GUIDELINES-UCSF HEALTH**INTRODUCTION**

The antimicrobial surgical prophylaxis guideline establishes evidence-based standards for surgical prophylaxis at UCSF Health. The protocol has been adapted from published consensus guidelines from the American Society of Health-System Pharmacists (ASHP), Society for Healthcare Epidemiology of America (SHEA), the Infectious Diseases Society of America (IDSA), Centers for Disease Control and Prevention (CDC), and the Surgical Infection Society (SIS) for use at UCSF with input from the Antimicrobial Stewardship Program, the Infectious Diseases Management Program, the Department of Anesthesiology, Allergy and Immunology, and the Surgical departments.

PRINCIPLES OF ANTIMICROBIAL SURGICAL PROPHYLAXIS

- This guideline is focused on clean and clean-contaminated surgeries
- Prophylaxis should target the most likely pathogens, taking into consideration type of surgery and local epidemiology
- Administer correctly—goal is for adequate tissue concentration at the time of infection risk
 - Administer within 60 minutes before the incision
 - For vancomycin and fluoroquinolones, start infusion 60-120 minutes prior to incision
- Give dose before the tourniquet goes up, if applicable
- Confirm timing of antibiotics with the surgeon at the Time-out or earlier since occasionally antibiotics need to be delayed until after culture
- In clean and clean-contaminated surgeries, discontinue antibiotics after the surgical incision is closed unless the patient has a documented or suspected infection. In pancreatic transplantation, continuation of antibiotics until duodenal cultures result is an exception.

Patients with existing infections

- The appropriate antibiotic to treat the underlying infection should be chosen on a case-by-case basis
- Continue the antibiotic to treat infection
- If spectrum of activity does not cover the usual organisms covered by routine prophylaxis for that type of case, add the routine prophylactic agent (discuss with ASP or infectious diseases if unclear)
- Ensure dose is given at appropriate time to achieve maximal tissue levels at time of incision (see above)
- Duration should be determined by the duration for the existing infection

Patients known to be colonized with methicillin-resistant *Staphylococcus aureus* (MRSA)

- Can consider addition of vancomycin to prophylaxis, especially if implant is being placed. Standard prophylaxis (e.g. cefazolin) should still be provided as this affords superior surgical site infection prevention for methicillin-sensitive *Staphylococcus aureus* (MSSA).
- If patients have a true contraindication to vancomycin (i.e. DRESS/SJS, anaphylaxis; these are very rare) can consider use of clindamycin in situations where vancomycin is indicated.

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025**Patients with recent history of resistant organisms**

- Data are limited
- Take into account:
 - Timing of infection
 - Location of infection
 - Prior treatment
 - Planned procedure
 - Organism

Patients with antibiotic allergy

- Take an antibiotic allergy history
 - Exact details of reaction, description of rash (if present)
 - Timing of reaction
 - Reason for antibiotic prescription
 - Other antibiotics received since then (also review the EMR to see whether the patient has received other antibiotics)
- Penicillin allergy: Patients with immediate type hypersensitivity to penicillins (including anaphylaxis, angioedema) can safely receive cefazolin and ceftriaxone. See [beta-lactam allergy guidance](#) for further details
 - Immediate-type hypersensitivity: Hives, angioedema, wheezing, anaphylaxis
 - Patients with immediate type hypersensitivity to penicillins, including anaphylaxis, can safely receive cefazolin and ceftriaxone (column 2)
 - Late reactions: Hemolytic anemia, thrombocytopenia, serum sickness, drug reaction with eosinophilia, Stevens Johnson syndrome (SJS)/Toxic epidermal necrolysis (TEN)
 - All beta-lactams should still be avoided, see alternative drugs (column 3)
- Cephalosporin allergy: consider allergy history to determine if rechallenge safe. See [beta-lactam allergy guidance](#) for further details.
 - If true cephalosporin allergy, use alternative drugs (column 3)
- When re-challenge or use of alternative β -lactam are likely safe:
 - Okay if patient had history of maculopapular rash (no hives, wheezing, anaphylaxis)
 - Okay if history of other drug intolerances like nausea

DOSING AND RE-DOSING INTERVALS

- In addition to the re-dosing intervals suggested below, consider immediate re-dosing in patients who have > 1.5 L of blood loss (>25ml/kg or > 30% blood volume loss for patients < 40kg) within a short time frame and those with severe burns.
 - Restart the re-dosing clock if this is done
 - Do not re-dose vancomycin or gentamicin for blood loss

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

COMMONLY USED IV DRUG DOSING AND ADMINISTRATION

ADULTS	Ampicillin-sulbactam [#]	Cefazolin	Ceftriaxone	Clindamycin	Gentamicin	Levofloxacin	Metronidazole	Piperacillin-tazobactam [#]	Vancomycin
Initial dose									
≥ 40 kg	3 g	40-120 kg: 2 g > 120 kg: 3 g	2 g	900 mg	5 mg/kg*	500 mg	500 mg	Standard: 3.375 g P aeruginosa: 4.5 g	40-80 kg: 1 g ≥ 80 kg: 1.5 g
< 40 kg adult	1.5 g	1 g	1 g	600 mg	2.5 mg/kg*	500 mg	500 mg	Standard: 3.375 g P aeruginosa: 4.5 g	750 mg
Admin	15 min IVPB	IVP over 3-5 mins	IVP over 3-5 mins	15 min IVPB	30 min IVPB Or 10 min IVPB	60 min IVPB	30 min IVPB	30 min IVPB	60 min/1 g IVPB
Intraoperative redosing interval, h	CrCl (>30): 2 CrCl (10-30): 6 CrCl (<10): Do not redose	CrCl (>30): 4 CrCl (10-30): 6 CrCl (<10): Do not redose	12	6	None	None	12	2	CrCl > 30: 12 CrCl ≤ 30: None
When to start standard dosing	After 3 doses	After 3 doses	After 2 doses	After 3 doses	Already standard	Already standard	Already standard	After 3 doses	Already standard
Standard dosing interval	q6h	q8h	q24h	q8h	q24h	q24h	Q12h	q8h	q12h
Compatibility issues	No	No	Ca ⁺⁺ containing solutions (incl LR) → possible precipitation 1. No Y-site 2. Flush line with normal saline or use alt solution	No	Propofol → Δ emulsion integrity 1. No Y-site 2. Flush line with normal saline	Propofol → free oil formation 1. No Y-site 2. Flush line with normal saline	Propofol → Δ fat globule size 1. No Y-site 2. Flush line with normal saline	Vecuronium → precipitation 1. No Y-site 2. Flush line with normal saline	No

[#]Note: Adult dosing is based on *total* drug product e.g. 3 g ampicillin-sulbactam = 2 g ampicillin + 1 g sulbactam

*Use ideal body weight, max dose = 400 mg

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

APPENDIX A. RECOMMENDED PROPHYLAXIS AGENTS BY ADULT OR PROCEDURE

	Recommended Agents	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
CARDIAC			
CABG	Cefazolin	Cefazolin	Vancomycin
Cardiac Device Insertion (pacemaker) and valves	Cefazolin	Cefazolin	Vancomycin
Ventricular Assist Device	MRSA nares (-): Cefazolin MRSA nares (+): Cefazolin + Vancomycin	MRSA nares (-): Cefazolin MRSA nares (+): Cefazolin + Vancomycin	Vancomycin
THORACIC			
Thoracotomy, Pneumonectomy, Lobectomy	Cefazolin	Cefazolin	Vancomycin
VATS	Cefazolin	Cefazolin	Vancomycin
GENERAL SURGERY			
GASTRODUODENAL—No suspected infection			
Entry into lumen of GI tract (bariatric)	Cefazolin	Cefazolin	Vancomycin + levofloxacin
Pancreatoduodenectomy (with or without stent)	Piperacillin/tazobactam	Ceftriaxone + metronidazole	Levofloxacin + metronidazole
Without entry into GI tract (anti-reflux)	Cefazolin	Cefazolin	Vancomycin + levofloxacin
BILIARY—No suspected infection			
Open-Biliary (no infection)	Cefazolin	Cefazolin	Levofloxacin + metronidazole
Laparoscopic- Biliary			
-Low Risk	None	None	None
-High Risk (no infection)	Cefazolin	Cefazolin	Levofloxacin + metronidazole
INTESTINE—No suspected infection			
Appendectomy (uncomplicated)	Cefazolin if > 8h from ceftriaxone; give metronidazole if not administered in the ED	Cefazolin if > 8h from ceftriaxone; give metronidazole if not administered in the ED	Levofloxacin + metronidazole
Nonobstructed small intestine	Cefazolin	Cefazolin	Clindamycin + levofloxacin
Hernia Repair	Cefazolin	Cefazolin	Vancomycin
Colorectal	Preop mechanical bowel prep + neomycin/metronidazole PO x 3 doses (metronidazole allergy: substitute erythromycin) Periop ceftriaxone + metronidazole	Preop mechanical bowel prep + neomycin/metronidazole PO x 3 doses (metronidazole allergy: substitute erythromycin) Periop ceftriaxone + metronidazole	Preop mechanical bowel prep + neomycin/metronidazole PO x 3 doses Periop levofloxacin + metronidazole

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

	Recommended Agents	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
HEAD & NECK			
Clean, including tympanostomy tube placement	None	None	None
Clean, neck dissection or prosthesis placement (except tympanostomy tube)	Cefazolin	Cefazolin	Clindamycin
FESS	Cefazolin	Cefazolin	Clindamycin
Tonsillectomy	None	None	None
Clean-contaminated except FESS and tonsillectomy	Ampicillin/sulbactam	Cefazolin + metronidazole	Clindamycin/levofloxacin
NEUROSURGERY			
Elective craniotomy & CSF Shunting-Primary	Cefazolin	Cefazolin	Vancomycin
Implantation of Intrathecal Pumps-Primary	Cefazolin	Cefazolin	Vancomycin
Revision craniotomy	Cefazolin + Vancomycin	Cefazolin + vancomycin	Vancomycin
Skull base with dural resection	Ceftriaxone (2 g IV q12H) + metronidazole (q8h)	Ceftriaxone (2g IV q12H) + metronidazole (q8h)	Aztreonam 2 g + metronidazole + vancomycin
ORTHOPEDIC and SPINE			
Clean operations of hand, knee, foot (without implantation of foreign material), ≤ 30 min	None	None	None
Clean operations of hand, knee, or foot (with foreign material), > 30 min	Cefazolin	Cefazolin	Vancomycin
Laminectomy or MIS primary spinal fusion	Cefazolin	Cefazolin	Vancomycin
Spinal fusion (primary): cervical or thoracic levels only	Cefazolin	Cefazolin	Vancomycin
Spinal fusion (primary): lumbar or sacral level involved	Cefazolin + gentamicin	Cefazolin + gentamicin	Vancomycin + gentamicin
Spinal fusion (revision): cervical or thoracic levels only	Cefazolin + vancomycin	Cefazolin	Vancomycin
Spinal fusion (revision): lumbar or sacral level involved	Cefazolin + vancomycin + gentamicin	Cefazolin + vancomycin + gentamicin	Vancomycin + gentamicin
Arthroplasty for femoral neck fracture	Cefazolin + (vancomycin if deemed high risk for drug-resistant organisms)	Cefazolin + (vancomycin if deemed high risk for drug-resistant organisms)	Vancomycin
Implantation of internal fixation devices (nails, screws, wires)	Cefazolin	Cefazolin	Vancomycin
Total joint replacement	Cefazolin	Cefazolin	Vancomycin
Total joint replacement- Revision	Cefazolin + vancomycin (cefazolin + clindamycin for a vancomycin allergy)	Cefazolin + vancomycin (cefazolin + clindamycin for a vancomycin allergy)	Vancomycin

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

	Recommended Agents	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
PLASTIC SURGERY			
Routine prophylaxis	Cefazolin	Cefazolin	Clindamycin
Approach through oral cavity	Ampicillin/sulbactam	Cefazolin + metronidazole	Clindamycin
TRANSPLANTATION			
Heart	Cefazolin	Cefazolin	Vancomycin
Heart with MRSA Risk Factors: Inpatient hospitalization >72 hrs in past 6 months Use of antibiotics in prior 48 hrs Open chest out of OR	Cefazolin + vancomycin	Cefazolin + vancomycin	Vancomycin + levofloxacin
Lung, Heart-Lung	Vancomycin + ceftriaxone	Vancomycin + ceftriaxone	Clindamycin
Kidney	Cefazolin	Cefazolin	Clindamycin
Liver	Piperacillin-tazobactam	Vancomycin + ceftriaxone	Vancomycin + Aztreonam
Pancreas & Kidney-Pancreas	Cefazolin +/- fluconazole (if high risk of fungal infection)	Cefazolin +/- fluconazole (if high risk of fungal infection)	Vancomycin + levofloxacin +/- fluconazole (if high risk of fungal infections)
UROLOGY			
Lower tract instrumentation with risk of infection (incl TURP)	Cephalexin (cefazolin if PO dose not given in preop)	Trimethoprim-sulfamethoxazole	Trimethoprim-sulfamethoxazole
Clean without entry into urinary tract	Cefazolin	Cefazolin	Clindamycin
Clean without entry into urinary tract + prosthesis	Cefazolin + gentamicin	Cefazolin + gentamicin	Vancomycin + gentamicin
Clean with entry into urinary tract	Cefazolin	Cefazolin	Clindamycin + gentamicin
Clean-contaminated	Ceftriaxone + metronidazole	Ceftriaxone + metronidazole	Levofloxacin + metronidazole
Complex stones/stents/upper tract instrumentation	Treat based on clinical judgement and culture data		
Upper tract instrumentation	Cephalexin (cefazolin if PO dose not given in preop)	Trimethoprim-sulfamethoxazole	Trimethoprim-sulfamethoxazole
Vaginal surgery	Cefazolin	Cefazolin	Clindamycin + gentamicin
VASCULAR	Cefazolin (+ vancomycin if MRSA colonization or graft/wound infection)	Cefazolin (+ vancomycin if MRSA colonization or graft/wound infection)	Vancomycin

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

APPENDIX B. RECOMMENDED PROPHYLAXIS AGENTS BY OB/GYN PROCEDURE

OB-GYN	Recommended Agents	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
Vaginal Delivery w/ manual placenta removal	Cefoxitin	Cefazolin + metronidazole	Clindamycin
Cesarean Delivery	Cefazolin	Cefazolin	Clindamycin + gentamicin
Caesarean Delivery contingencies	<p><u>MRSA colonized</u>: consider vancomycin x 1, in addition to above</p> <p><u>Ruptured membranes or labor at time of C-section</u>: Add Azithromycin 500 mg IV x 1</p> <p><u>Vaginal hand</u>: Standard prophylaxis as above</p> <p><u>Stat C-section</u>: Standard prophylaxis as above</p>		
Vaginal packing	No targeted prophylaxis unless another indication for prophylaxis exists		
Uterine balloon	Cefoxitin	Cefazolin + metronidazole	Clindamycin
Laceration repair (3 rd or 4 th degree only)	Cefoxitin 2g IV	Cefazolin + metronidazole	Clindamycin
Hysterectomy (vaginal or abdominal), urogynecology procedures (+/- mesh)	Cefazolin +/- metronidazole	Cefazolin +/- metronidazole	Clindamycin + gentamicin
Hysteroscopy, laparoscopy, laparotomy	None		
Hysterosalpingogram, endometrial biopsy	Doxycycline 100 mg po twice daily x 5 days if hx of PID or dilated fallopian tubes		
Medical abortion	None		
Induced abortion/D+E/ D+C/manual uterine inspection (including Laminaria placement)	Doxycycline 200 mg po prior to procedure		
With fetal demise (any gest age)	Add metronidazole 500 mg q8H x 2 doses		
Ultrasound (USIC), history-indicated cerclage (HIC), or laparoscopic-assisted abdominal cerclage	None		
Physical exam-indicated cerclage (PEIC)	Consider cefazolin + indomethacin	Consider cefazolin + indomethacin	Consider clindamycin + indomethacin
Transabdominal cerclage (TAC)	Cefazolin	Cefazolin	Clindamycin
<u>Low risk</u> : no antibiotics required			

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

OB-GYN	Recommended Agents	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
Cystoscopy alone	<u>High risk:</u> TMP/SMX 160/800 DS (+ urine culture, pre-op catheter, prosthetic material)		
Cystoscopy with manipulation or upper tract instrumentation (e.g. ureteroscopy)	TMP/SMX 160/800 DS x1		
Cystoscopy with open surgery	Cefazolin	Cefazolin	Clindamycin + gentamicin
Risk of colorectal injuries	Preoperative mechanical bowel prep + neomycin/metronidazole PO x 3 doses Perioperative ceftriaxone + metronidazole	Preoperative mechanical bowel prep + neomycin/metronidazole PO x 3 doses Perioperative ceftriaxone + metronidazole	Preoperative mechanical bowel prep + neomycin/metronidazole PO x 3 doses Perioperative levofloxacin + metronidazole
PUBS/IUT/RFA/Thoracoamniotic shunt/vesicoamniotic shunt/amnioreduction/ECV/laser ablation in TTTS, FETO (fetal endoscopic balloon occlusion), myelomeningocele repair/open fetal surgery	Cefazolin	Cefazolin	Clindamycin

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

APPENDIX C. RECOMMENDED PROPHYLAXIS AGENTS BY IR PROCEDURE

	Recommended agent	Immediate reactions to penicillin (including anaphylaxis, hives, angioedema)	Cephalosporin allergy or delayed severe penicillin reactions (Type II-IV: SJS/TEN, DRESS, serum sickness)
Angiography, angioplasty, thrombolysis, arterial closure device placement, stent placement, superficial venous insufficiency treatment, IVC filter placement	None	None	N/A
Central venous access	None	None	None
Chemoembolization, ablation, embolization, or radioembolization where sphincter of Oddi is not intact (eg, history of biliary reconstructive surgery)	Moxifloxacin 400 mg daily x 3 days prior and 11 days post-procedure		
Embolization, ablation, radioembolization, and chemoembolization of liver (if intent to create infarction or high likelihood of infarction)	Ampicillin/sulbactam	Cefazolin + metronidazole	Levofloxacin
Endograft placement	None	None	None
Gastrostomy tube placement	Cefazolin	Cefazolin	Levofloxacin
Percutaneous abscess drainage, on antibiotics	Continue directed therapy for the existing infection		
Percutaneous abscess drainage, not on antibiotics	Discuss with primary service whether antibiotics should be given (if active signs of infection) or held until after cultures obtained		
Percutaneous nephrostomy tube placement or change	Cefazolin	Cefazolin	Levofloxacin
Renal ablation, embolization	Cefazolin	Cefazolin	Levofloxacin
Uterine Artery Embolization (UAE)	Cefazolin	Cefazolin	Levofloxacin
TIPS creation	Cefazolin	Cefazolin	Levofloxacin
Tube check/change (hepatic, biliary)	Cefazolin	Cefazolin	Levofloxacin

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

REFERENCES

ACOG Practice Bulletin 104. Antibiotic prophylaxis for gynecologic procedures. May 2009 (reaffirmed 2016)

Ancef [package insert]. Triangle Park, NC: GlaxoSmithKline; 2004

Allegranzi B, et al. Surgical site infections 2: New WHO recommendations on intraoperative and postoperative measures for surgical site infection prevention: an evidence-based global perspective. *Lancet Infect Dis* 2016(12): e276-e287

Ariyan S, Martin J, Lal A, Cheng D, Borah GL, Chung KC, Conly J, Havlik R, Lee WPA, McGrath MH, Pribaz J, Young VL: Antibiotic Prophylaxis for Preventing Surgical-Site Infection in Plastic Surgery: an Evidence-Based Consensus Conference Statement from the American Association of Plastic Surgeons. *Plast Reconstr Surg* 135(6): 1723-1739, 2015

Berrios-Torres SI, Umscheid CA, Bratzler DW, et al. Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. *JAMA Surg* Published online May 3, 2017. doi:10.1001/jamasurg.2017.0904

Blumenthal, K. G., Ryan, E. E., Li, Y., Lee, H., Kuhlen, J. L., & Shenoy, E. S. (2018). The Impact of a Reported Penicillin Allergy on Surgical Site Infection Risk. *Clinical Infectious Diseases*, 66(3), 329–336. <https://doi.org/10.1093/CID/CIX794>

Bratzler DW, Dellinger EP, Olsen KM, et al. American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm*. 2013;70(3):195-283

Cefoxitin [package insert]. Weston, FL: Apotex Corp; 2015.

Cleocin [package insert]. New York, NY: Pfizer; 2005

Kuruvilla, M., Sexton, M. E., Wiley, Z., Langfitt, T., Lynde, G., & Wolf, F. (2020). A streamlined approach to optimize perioperative antibiotic prophylaxis in the setting of penicillin allergy labels. *Journal of Allergy and Clinical Immunology*, 145(2), AB55. <https://doi.org/10.1016/j.jaci.2019.12.696>

Nelson RL, Gladman E, Barbateskovic M. Antimicrobial prophylaxis for colorectal surgery. *Cochrane Database Syst Rev*. 2014;(5):CD001181

Venkatesan AM, Kundu S, Sachs D, et al. Practice guideline for adult antibiotic prophylaxis during vascular and interventional radiology procedures. *J Vasc Interv Radiol* 2010; 21: 1611-1630

Unasyn [package insert]. New York, NY: Pfizer Inc; 2007

Owner, UCSF Antimicrobial Stewardship Program (Will Simmons MD)

Adult antimicrobial surgical prophylaxis guidelines
Approved 08.2025, Last update 08.2025

Zosyn [package insert]. New York, NY: Pfizer; 2012