

Sensitivity Patterns of NON-URINE Isolates (First isolate per patient per organism)

San Francisco VA Medical Center

January-December 2021

Confidential: For official use by San Francisco VA Medical Center Employees and Students only

<u>Percent Sensitive</u>						<u>Percent Sensitive</u>			
ORGANISM	TOTAL #ISO*	AMP	TZP	ETP	CIP	SXT	CRO	FEP	GEN
<i>Enterobacter cloacae</i>	30	NA	NA	100	100	93	NA	100	100
<i>Escherichia coli</i> ESBL (10%)	69	55	97	100	71	74	84	90	87
<i>Klebsiella pneumoniae</i> ESBL (12%)	34	NA	94	100	94	85	88	88	97
<i>Proteus mirabilis</i>	28	75	100	100	79	86	75	75	89
<i>Pseudomonas aeruginosa</i>	56	NA	91	NA	93	NA	NA	96	98
		AMP	DAP	LZD	VAN	OXA	SXT	CLI	TET
<i>Enterococcus faecalis</i> VRE (0%)	49	97	100	100	100	NA	NA	NA	NA
<i>Staphylococcus aureus</i>	222	NA	100	100	100	71	97	81	91
MRSA (29%)	64	NA	100	100	100	0	92	67	77
MSSA (71%)	158	NA	100	100	100	100	100	87	95
<i>Staphylococcus coag neg</i>	104	NA	100	100	93	48	57	57	86
Methicillin Resistant (52%)									

NA = not available

AMP - ampicillin, TZP piperacillin/tazobactam, ETP - ertapenem, CIP - ciprofloxacin, SXT - trimethoprim/sulfamethoxazole

CRO - ceftriaxone, FEP - cefepime, GEN - gentamicin, VAN - vancomycin, OXA - oxacillin, CLI - clindamycin, TET - tetracycline

*Statistical validity of % susceptible is decreased if fewer than 30 isolates are tested.