

Sensitivity Patterns of URINE Isolates (First isolate per patient per organism)

San Francisco VA Medical Center

January-December 2015

Confidential: For official use by San Francisco VA Medical Center Employees and Students only

ORGANISM	TOTAL #ISO*	Percent Sensitive							
		AMP	AUG	CEPH	CIP	GEN	TOB	NITR	T/S
<i>Enterobacter cloacae</i>	22	NA	NA	NA	100	100	100	91	91
<i>Escherichia coli</i> †	305	49	90	78	75	90	97	98	72
<i>Klebsiella oxytoca</i> †	30	0	100	63	97	100	100	100	97
<i>Klebsiella pneumoniae</i> †	90	0	99	96	99	97	97	91	89
<i>Morganella morganii</i>	23	0	0	0	83	83	90	0	65
<i>Proteus mirabilis</i>	61	70	100	56	75	89	100	0	74
<i>Pseudomonas aeruginosa</i>	54	NA	NA	NA	91	94	95	NA	NA
					OXAC	VANC	TCN	NITR	T/S
<i>Staphylococcus aureus</i>	42				67	100	98	100	90
<i>Staphylococcus coag neg</i>	165				56	99	82	100	69

NA = not available

AMP - ampicillin, AUG - amoxicillin/clavulanate, CEPH - cefazolin, CIP - ciprofloxacin, GEN - gentamicin, NITR - nitrofurantoin,

TCN - tetracycline, T/S - trimethoprim/sulfamethoxazole, OXAC - oxacillin, VANC - vancomycin

*Statistical validity of % susceptible is decreased if fewer than 30 isolates are tested.

10.5% (41/390) of all enterococcal isolates were vancomycin-resistant

†Extended-Spectrum Beta-Lactamase (ESBL) Positive - *E. coli* - 7%; *K. oxytoca* - 0%; *K. pneumoniae* - 1%